

Lattice four-dimensional $\mathcal{N}=4$ SYM is practical

Joshua W. Elliott^a, Joel Giedt^b and Guy D. Moore^a

^a *Physics Department, McGill University, 3600 rue University, Montréal, QC H3A 2T8, Canada*

^b *Department of Physics, Applied Physics, and Astronomy,
Rensselaer Polytechnic Institute, 110 Eighth Street, Troy, New York 12180-3590 USA*

(Dated: 30 May 2008)

We show that nonperturbative lattice studies of four-dimensional $\mathcal{N}=4$ Super-Yang-Mills are within reach. We use Ginsparg-Wilson fermions to avoid gluino masses and an exact implementation of the (chiral) R -symmetry, which greatly limits the number of counterterms that must be fine-tuned. Only bosonic operators require fine tuning, so all tunings can be done “offline” by a Ferrenberg-Swendsen type reweighting. We show what measurables can be used to perform the tuning.

PACS numbers:

It is often assumed that the study of supersymmetry (SUSY) on the lattice will be difficult in four dimensions because multiple relevant/marginal operators that violate SUSY are allowed by lattice symmetries. A notable exception is pure $\mathcal{N}=1$ super-Yang-Mills (SYM). In that case, lattice chiral symmetry in the form of Ginsparg-Wilson (GW) fermions prevents additive renormalization of the gluino mass in the continuum limit, and hence the only SUSY-violating operator is forbidden in that limit by setting the bare mass to zero [1] $\mathcal{N}=4$ SYM has scalars and Yukawa couplings, and so various masses and couplings will receive divergent corrections in the continuum limit and must be (nonperturbatively) fine-tuned away. Hence lattice $\mathcal{N}=4$ SYM has always seemed impractical by this fine-tuning approach.[13]

We show here that this is not the case. Using GW fermions the four gluinos can be kept massless and the $SU(4)_R$ symmetry in the lattice theory can be preserved, which greatly restricts the form of the renormalizations. This leaves a scalar mass, two quartic couplings, and a Yukawa coupling to tune. The Yukawa coupling can be tuned by rescaling the scalar kinetic term (see below); so all tunings can be done by adjusting bosonic terms in the action. This allows the tunings to be done by the “Ferrenberg-Swendsen method” [2, 3], exploring a wide swath of coupling constant space “offline” from the results of a single Monte-Carlo simulation. The parameter range available with good statistics can be enlarged using multicanonical techniques [4, 5]. Thus we arrive at the encouraging result that all fine-tuning can be performed through an “offline” analysis, *i.e.*, new simulations and fermion matrix inversions are not required.

The continuum field content is $SU(N_c)$ YM theory with an $SU(4)_R$ internal symmetry; there are four Majorana fermions whose left handed components transform in the fundamental $\mathbf{4}$ representation of $SU(4)_R$ and 6 real scalars in the antisymmetric tensor representation. The six real scalars will be expressed with a single index $\phi_m, m=1\dots 6$, or composed into $SU(4)_R$ Weyl matrices: $\phi_{ij}=\phi_m\hat{\sigma}_{m,ij}$ and $\phi^{ij}=\phi_m\hat{\sigma}_m^{ij}$, where $\hat{\sigma}$ ’s are just $SU(4)_R$ Clebsch-Gordon coefficients involved in $\mathbf{4}^*\otimes\mathbf{6}\otimes\mathbf{4}$. The

continuum action is

$$S = \frac{1}{g^2} \text{Tr} \left\{ \frac{1}{2} F^2 + |D_\mu \phi_m|^2 + \bar{\psi}_i \not{D} \psi_i + \sqrt{2} \bar{\psi}_i (\phi^{ij} P_L - (\phi^{ij})^* P_R) \psi_j + [\phi_m, \phi_n][\phi_m, \phi_n] \right\}. \quad (1)$$

The $SU(4) \simeq SO(6)$ preserving bosonic lattice action is a trivial transcription from the continuum. Of course we must allow for generic coefficients and non-SUSY terms, so that the SUSY-restoring counterterms can be tuned. In our case these are entirely scalar terms.

It has been argued that naive lattice Yukawa terms lead to inconsistencies in either the chiral or Majorana projections (depending on how the Yukawas are transcribed to the lattice) [6], so the fermionic implementation is more subtle. Following Lüscher [7] and Kikukawa and Suzuki [8], we introduce a set of auxiliary fermionic fields Ψ , with lattice fermionic action

$$S_F = a^4 \text{Tr} \left\{ \bar{\psi}_i D_{\text{GW}} \psi_i - a^{-1} \bar{\Psi}_i \Psi_i + y \sqrt{2} (\bar{\psi} + \bar{\Psi})_i (\phi^{ij} P_L - (\phi^{ij})^* P_R) (\psi + \Psi)_j \right\}. \quad (2)$$

This action possesses an exact $SU(4)_R$ symmetry, with the scalars transforming as in the continuum and the fermions transforming according to

$$\begin{aligned} \delta \psi / i\varepsilon &= (T \hat{P}_L - T^* \hat{P}_R) \psi, \\ \delta \Psi / i\varepsilon &= (T + T^*) \gamma_5 D_{\text{GW}} \psi + (T P_L - T^* P_R) \Psi, \\ \delta \bar{\psi} / i\varepsilon &= \bar{\psi} (T^* P_L - T P_R) + \bar{\Psi} (T + T^*) \gamma_5, \\ \delta \bar{\Psi} / i\varepsilon &= -\bar{\Psi} (T P_L - T^* P_R). \end{aligned} \quad (3)$$

Here $\hat{P}_{L/R} \equiv \frac{1}{2}(1 \pm \hat{\gamma}_5) = \frac{1}{2}(1 \pm \gamma_5(1 - 2D_{\text{GW}}))$ are the lattice modified chiral projection operators, T is the generator of $SU(4)_R$ in the fundamental ($\mathbf{4}$) and we have suppressed the $SU(4)_R$ indices. Hence $(\psi + \Psi)$ and $(\bar{\psi} + \bar{\Psi})$ transform like the continuum $\psi, \bar{\psi}$ fields.

This auxiliary field method preserves the R -symmetry exactly and keeps the Yukawa terms ultralocal. It is also consistent with the Majorana decomposition, so the fermionic determinant is an exact square; taking its square root to implement the Majorana nature of the fermions retains locality. The cost is the introduction of

an extra fermionic excitation Ψ , which is however nondynamical with $\mathcal{O}(a^{-1})$ mass, so it decouples from the theory in the continuum limit. Note however that in building lattice operators which contain fermions, we must always use the combination $\psi + \Psi$, since this is the combination with correct R -symmetry transformations.

We are interested in tuning the lattice action such that the effective infrared description is $\mathcal{N}=4$ SYM. Generically there is a nontrivial matching between lattice and effective IR theories and all relevant or marginal terms consistent with lattice symmetries will appear in the infrared, except at special points in bare parameter space. We can arrive at the desired special point (*i.e.*, $\mathcal{N}=4$ SYM) by introducing the SUSY-violating operators into the bare action and fine-tuning counterterms. These counterterms fall into three categories: a scalar mass term, a Yukawa term, and two or four scalar quartic terms, depending on the number of colors for the gauge group, restricted here to $SU(N_c)$. In particular, if $N_c \leq 3$ then $SU(4)_R$ symmetry allows only two unique quartic terms: $(\text{Tr} \phi_m \phi_m)^2$ and $(\text{Tr} \phi_m \phi_n)^2$ (trace over gauge indices, R -indices explicit). For $N_c > 3$ the quartic invariants $\text{Tr} \phi_m \phi_m \phi_n \phi_n$ and $\text{Tr} \phi_m \phi_n \phi_m \phi_n$ are also independent (for $N_c \leq 3$ they are linearly dependent on the first two).

Rescaling the Yukawa term, schematically $y \bar{\psi} \phi \psi$, can be accomplished through a rescaling of the scalar kinetic term. Consider replacing $y \bar{\psi} \phi \psi + |D\phi|^2$ with $y \bar{\psi} \phi \psi + Z |D\phi|^2$. In terms of the canonically normalized scalar field this is $(y/\sqrt{Z}) \bar{\psi} \phi \psi + |D\phi|^2$. Therefore we include a counterterm for the kinetic term rather than for the Yukawa. (Note that the rescaling of the scalar kinetic term also rescales the scalar potential; but there are distinct counterterms to undo this rescaling separately).

Suppose we perform a Monte Carlo simulation at one value m_1 of the scalar mass m , so that the configurations sample the distribution determined by the action $S = S_{m=0} + \frac{1}{2} \int m_1^2 \phi^2$. Following the “Ferrenberg-Swendsen reweighting” method [2, 3] one can use the following “reweighting identity” to compute the expectation value of an operator O for the distribution with a mass m_2 :

$$\langle O \rangle = \sum_c O_c e^{-\frac{1}{2}(m_2^2 - m_1^2) \int \phi_c^2} / \sum_c e^{-\frac{1}{2}(m_2^2 - m_1^2) \int \phi_c^2}. \quad (4)$$

There is a limited regime of utility to this technique, due to the so-called “overlap problem.” For instance, if the exponential in (4) is large where the simulated distribution has little weight, a finite sampling will have large errors. The mismatch of the distributions gets worse as the number of lattice sites increases, because the exponent is extensive.

A way to ameliorate the overlap problem, which has been found to work in other contexts, is “multicanonical reweighting” [4]. One replaces S with $S + W[O_1, O_2, \dots]$, where $W[O_1, O_2, \dots]$ is a carefully chosen function of

some small set of observables (in our case W will be a function of $\int \phi^2$, the distinct $\int \phi^4$ ’s, and $\int (D\phi)^2$). The expectation value of an observable in the distribution corresponding to S is:

$$\langle O \rangle = \sum_c O_c e^{W[O_1^c, \dots]} / \sum_c e^{W[O_1^c, \dots]} \quad (5)$$

$W[\dots]$ produces a weighted average over a continuum of canonical ensembles, some of which will have a good overlap with the distribution that one is reweighting to. The challenge is to design a W such that sampling is flattened over the range of observables one is interested in.

Two approaches to engineering a good function W exist: (1) a bootstrap method that iterates between Monte Carlo simulation and adjusting W , and (2) optimizing W w.r.t. its parameters, in a small volume, and then using step-scaling to extrapolate to a good estimate for W in larger volumes. For instance one can start with 4^4 and 6^4 volumes, where statistics accumulate rapidly and unweighted simulations still cover broad parameter ranges.

In our case the reweighting function W will depend on the four bosonic contributions to the action, $\int \phi^2$, $\int (D\phi)^2$, $\int \phi_1^4$ and $\int \phi_2^4$ (the two quadratic and two independent quartic operators you can form from the scalars, integrated over space). By sampling with the weight

$$\rho = \mathcal{D}[A, \phi] \det[D] e^{-S[A, \phi; m_0^2, \dots]} e^{-W[\int \phi^2, \dots]}, \quad (6)$$

and you can reproduce the ensemble at some particular set of values for m^2 , Z_ϕ , λ and λ' , via

$$Z = \sum_c e^{+W[\dots]} e^{-[(m^2 - m_0^2) \phi_c^2 + (Z_\phi - Z_\phi^0) (D\phi_c)^2 + \dots]}, \quad (7)$$

with W chosen so that the sample has a reasonable number of configurations for all values of $\int \phi^2$, $\int (D\phi)^2$, $\int \phi^4$ and $\int (\phi^2)^2$ within some interesting range.

Now, define the gauge invariant effective potential in finite volume as follows:

$$e^{-\Omega V[A^2]} = \sum_c e^{+W[\dots]} e^{-[(m^2 - m_0^2) \phi_c^2 \dots]} \times \delta(A^2 - \frac{\phi_c^2}{\Omega}) \quad (8)$$

with Ω the 4-volume (so that A^2 represents the mean value of the squared scalar field). It is easy to check that varying m^2 changes $V[A^2]$ by adding a linear component. Therefore, measuring $V[A^2]$ immediately determines how $\langle \phi^2 \rangle$ and $F = \ln Z$ vary as a function of m^2 . One easily generalizes to the effective potential as a function of all four bosonic operators, which gives a quick way to explore the effect of varying parameters.

It remains to specify how to tune the parameters. Consider $\mathcal{N}=4$ SYM with an added $a^2 (\text{Tr} \phi_m \phi_m)^3$ term in the potential, but deformed by mass and quartic interactions. The SUSY point is a second order phase transition point; for negative quartic deformation there is a first order transition as m^2 is varied (the system jumps from a

massive state about the origin to a massive state with a large VEV because the quartic term bends down). For positive quartic deformation there is no phase transition. The optimal value of m^2 (fixing other parameters) is the point of maximal susceptibility $\langle(\int_x \phi^2)^2\rangle - (\langle\int_x \phi^2\rangle)^2$. Determining this point in finite volume leads to $\mathcal{O}(1/L^2)$ errors in the determined value of m^2 , which can be improved by scaling over multiple volumes. Finding the flat quartic term which gives second-order behavior should also be possible; it has been successfully achieved in the context of the electroweak phase transition [9]. Therefore it should be possible to use the phase diagram to tune at least two parameters. Note that we needed to add a lattice-size ϕ^6 term; this is harmless but it raises the issue that the flat direction actually means that unbroken $\mathcal{N}=4$ SYM is not well behaved in finite volume; the moduli are not fixed and the partition function diverges because of the integral over the infinite moduli space. Therefore it will always be necessary to break SUSY somehow. We advocate doing so via twisted boundary conditions; for instance, instead of periodic boundary conditions we can add a rotation by angle Θ to all fermionic fields in one direction. The choice $\Theta=\pi$ is the maximal global breaking of SUSY and corresponds to treating the thermal ensemble; intermediate values of Θ break SUSY by smaller amounts. This lifts the moduli degeneracy without any *local* SUSY breaking; the effects of Θ are only visible in correlations at the scale of the lattice size, which is anyway contaminated by being in finite volume.

For the case $N_c > 3$ the effective potential should show multiple flat directions in the space of quartic operators; only one quartic direction (some linear combination of the input quartics, due to mixing) should rise steeply. Therefore we expect it should be possible to tune the “extra” quartic operators in the case $N_c > 3$, leaving only one quartic and the Yukawa coupling/wave function to tune.

If SUSY is exact then the (R -symmetry 4) supercurrent $S_{\mu,i}$ is conserved, so $\langle\partial^\mu S_{\mu,i}(x)\mathcal{O}(y)\rangle$ vanishes at $x\neq y$ for all local operators $\mathcal{O}$. We can use this property to measure whether we are at the SUSY point in parameter space, and therefore to tune parameters to find the SUSY point. The technique has been pioneered in $\mathcal{N}=1$ SUSY with Wilson fermions by the DESY-Münster group [10]; here we discuss the extension to $\mathcal{N}=4$ SYM.

The supercurrent $S_{\mu,i}$ is a linear combination of three dimension-7/2 operators. It is easy to find lattice operators which reproduce these continuum operators, at tree level and with contamination from higher dimension operators. The choices are not unique and at the non-perturbative level each lattice operator will mix with all continuum operators in the same symmetry channel. Different choices of lattice operator will reproduce the continuum operator with different normalization, mixings, and $\mathcal{O}(a)$ suppressed higher dimension contamination. Hence we express the operators $\mathcal{O}_{\mu,i}$ in a continuum lan-

guage, and leave the particulars of lattice transcription (which amounts to various “improvements” w.r.t. $\mathcal{O}(a)$ discretization errors) for detailed studies. Our intention here is to lay out the methodology.

In their analysis of the $\mathcal{N}=1$ SYM case, the DESY-Münster group found two dimension-7/2 operators, the supercurrent S_μ and another fermionic current T_μ . These mix in the lattice-continuum matching and so one must write down two lattice operators with undetermined coefficients in order to find something which corresponds purely to S_μ (plus $\mathcal{O}(a)$ dimension-9/2 contamination). In the present $\mathcal{N}=4$ case there are 5 dimension-7/2 operators which we will name $\mathcal{O}_{\mu,i}^{1\dots 5}$, and the renormalized $\mathcal{N}=4$ supercurrent will, in all generality, take the form:

$$\begin{aligned} S_{\mu,i}^{\text{ren.}} = & \{Z_1 \tfrac{1}{2} F_A \cdot \sigma \delta_{ij} + Z_2 \sqrt{2} \mathcal{P}(\phi_{ij} P_L + \phi^{ij} P_R)_A \\ & - Z_3 f^{ABC} (\phi_{ik}^B \phi_C^{kj} P_R + \phi_B^{ik} \phi_{kj}^C P_L)\} \gamma_\mu \psi_{jA} \\ & + \{Z_4 \gamma_\nu F_{\mu\nu}^A \delta_{ij} - Z_5 \sqrt{2} D_\mu (\phi^{ij} P_L + \phi_{ij} P_R)_A\} \psi_{jA} + \mathcal{O}(a) \\ \equiv & Z_n \mathcal{O}_{\mu,i}^n + \mathcal{O}(a), \end{aligned} \quad (9)$$

where the terms on the righthand side are bare (lattice) operators. Note that: (1) at tree level the supercurrent corresponds to $Z_1=Z_2=Z_3=1$ and $Z_4=Z_5=0$; (2) the renormalization constants Z_n are universal w.r.t. the index i due to the $SU(4)_R$ symmetry preserved by the lattice.

We can tune to the SUSY point by varying parameters to force correlation functions of this lattice-implemented $\partial_\mu S_{\mu,i}$ to vanish up to $\mathcal{O}(a)$ corrections. Specifically, to tune two parameters we need to choose 6 operators $\mathcal{O}_{\mu,i}^n$ in the same symmetry channel as $S_{\mu,i}$ (otherwise the correlation function vanishes automatically). The natural choice is $\mathcal{O}_{\mu,i}^{1\dots 5}$ plus one dimension-9/2 operator $\mathcal{O}_{\mu,i}^6$. One then measures the matrix of correlation functions

$$M^{mn}(t) \equiv \int d^3\vec{x} \langle \mathcal{O}_{0,i}^{m\dagger}(t, \vec{x}) \mathcal{O}_{0,i}^n(0, 0) \rangle \quad (10)$$

whose t derivative is the correlation function between $\partial_\mu \mathcal{O}_{\mu,i}^m$ and $\mathcal{O}^n(0, i)$ at vanishing spatial momentum. Since the operators involved are dimension-7/2 we generically expect the elements of $M^{mn}(t)$ to decay as t^{-7} . At the SUSY point and for the right choices of Z_m , $Z_m M^{mn}$ decays as at^{-8} for all n . We can fix the undetermined ratios $Z_{2\dots 5}/Z_1$ by enforcing that this holds for $n=1\dots 4$. Forcing that it hold for $n=5, 6$ gives two conditions which can be used to check whether we are at the SUSY point—tuning to the SUSY point is tuning for $Z_m M^{m5} \sim at^{-8}$ and $Z_m M^{m6} \sim at^{-8}$. Actually since one of the operators is dimension-9/2 we must force one linear combination $Z_m M^{mn} c_n$ to vanish as $a^2 t^{-9}$.

We do not see an obstacle to using this procedure to tune more parameters, if it proves too difficult to tune some of the vanishing quartic couplings via the potential method. Therefore in principle the tuning to the SUSY point can be done by any mixture of the Ward identity method and the effective potential method.

We have seen that for $N_c=2,3$ colors, there are four fine-tunings in the action. For $N_c>3$ colors there are six. In addition, one must fix the four relative renormalization constants in the supercurrent. All but one of the scalar potential counterterms can be fixed by matching the effective potential, as determined by the multiccanonical simulation, to the target theory scalar potential $\text{Tr}[\phi_m, \phi_n][\phi_m, \phi_n]$. The overall strength of this term cannot be determined from the effective potential, because it will be expressed in terms of the bare operators in our approach.

This leaves just six fine-tunings for all number of colors N_c : one fine-tuning of the bare kinetic coefficient for the scalar, one overall scalar potential coefficient, and the four relative supercurrent coefficients. Thus a total of six Ward identities must be measured well enough to distinguish their simultaneous minimum w.r.t. $Z_\phi, Z_1/Z_2, \dots, Z_4/Z_5$.

Discussion

By preserving the $SU(4)_R$ symmetry of the target theory, the number of counterterms that must be fine-tuned is greatly reduced. This can be done by implementing GW fermions, with the chirality of the Yukawa couplings implemented with auxiliary fermions, extending the method of [8]. Because counterterm fine-tuning can be isolated to the purely bosonic sector, it can all be done off-line, *i.e.*, without the expense of fermion matrix inversions (the bottleneck for all dynamical fermion simulations). This is a great advantage, because a very large number of points in the bare action parameter space will have to be scanned in order to find the $\mathcal{N}=4$ SYM point. Finally, we have explained how the overlap problem can be alleviated by taking a multiccanonical approach, flattening the distributions that will be scanned over.

The main limitation to this method is that, since $\mathcal{N}=4$ SYM is conformal, the continuum limit is not a weak coupling limit. Our proposal should work at weak coupling, where one knows that the infrared description will be in terms of the same degrees of freedom as one puts on the lattice. But there is no guarantee that one can find lattice parameters which correspond to strongly coupled continuum theories.

The principal challenge is that the method requires GW fermions, which are numerically expensive—especially in a theory such as this one, with massless particles and the corresponding critical slowing down. It will be a challenge to generate enough configurations to measure quantities with sufficient accuracy to determine the SUSY point. Fermions are necessarily involved in the correlation functions of the supercurrent, so storage of propagators during the simulations will be essential to performing the fine-tuning w.r.t. Ward identities. The

storage and computing resources that will be required will be substantial, but we believe that the exploratory studies that need to be done can be performed in the near term. For instance, one of the authors (JG) has access to the Computational Center for Nanotechnology Innovation (CCNI) at Rensselaer. Indeed, it is currently being used by Giedt and collaborators for $\mathcal{N}=1$ SYM simulations at a sustained actual compute rate of 1 Tflop/s, precisely the sort of resource that would be required to perform exploratory studies of this proposal. Obviously early stages of such work will be very much technical studies of the lattice theory. Continuum results will take much longer. Nevertheless, the beginnings of first principles nonperturbative study of $\mathcal{N}=4$ SYM are not so far off, we believe, if the current proposal is pursued with some dedication and adequate resources. We hope to report on further progress in that direction in the near future.

Acknowledgements

This work was supported in part by the Natural Sciences and Engineering Research Council of Canada.

-
- [1] G. Curci and G. Veneziano, Nucl. Phys. **B292**, 555 (1987). N. Maru and J. Nishimura, Int. J. Mod. Phys. **A13**, 2841 (1998). H. Neuberger, Phys. Rev. **D57**, 5417 (1998). J. Nishimura, Phys. Lett. **B406**, 215 (1997). D. B. Kaplan and M. Schmaltz, Chin. J. Phys. **38**, 543 (2000). I. Campos et al. (DESY-Munster), Eur. Phys. J. **C11**, 507 (1999). G. T. Fleming, J. B. Kogut, and P. M. Vranas, Phys. Rev. **D64**, 034510 (2001). R. Kirchner, S. Luckmann, I. Montvay, K. Spanderen, and J. Westphalen (DESY-Munster), Nucl. Phys. Proc. Suppl. **73**, 828 (1999). A. Donini, M. Guagnelli, P. Hernandez, and A. Vladikas, Nucl. Phys. **B523**, 529 (1998).
 - [2] M. Falcioni, E. Marinari, M. L. Paciello, G. Parisi, and B. Taglienti, Phys. Lett. **B108**, 331 (1982).
 - [3] A. M. Ferrenberg and R. H. Swendsen, Phys. Rev. Lett. **61**, 2635 (1988).
 - [4] B. Baumann, Nucl. Phys. **B285**, 391 (1987).
 - [5] K. Kajantie, M. Laine, K. Rummukainen, and M. E. Shaposhnikov, Nucl. Phys. **B466**, 189 (1996).
 - [6] K. Fujikawa, M. Ishibashi, and H. Suzuki, Phys. Lett. **B538**, 197 (2002).
 - [7] M. Luscher, Phys. Lett. **B428**, 342 (1998).
 - [8] Y. Kikukawa and H. Suzuki, JHEP **02**, 012 (2005).
 - [9] K. Rummukainen et al., Nucl. Phys. **B532**, 283 (1998), hep-lat/9805013.
 - [10] F. Farchioni et al. (DESY-Munster-Roma), Eur. Phys. J. **C23**, 719 (2002).
 - [11] S. Catterall, JHEP **06**, 027 (2005).
 - [12] D. B. Kaplan and M. Unsal, JHEP **09**, 042 (2005).
 - [13] Other approaches recently suggested include those of Refs. [11, 12], involving “orbifold” or “twisted SUSY” lattices.